



Anaphylaxis Policy

Mornington Park Primary School 5040

POLICY DATE: January 2026

STATUS: Approved by School Council

REVIEW: January 2027

“Victorian government schools are child safe environments. Our schools actively promote the safety and wellbeing of all students, and all school staff are committed to protecting students from abuse or harm in the school environment in accordance with their legal obligations including the Child Safe Standards. Mornington Park Primary School’s (The school’s) Child Safety Code is available on the school’s website.”

PURPOSE

To explain to Mornington Park Primary School, parents, carers, staff and students the processes and procedures in place to support students diagnosed as being at risk of suffering from anaphylaxis. This policy also ensures that Mornington Park Primary School is compliant with Ministerial Order 706 and the Department’s guidelines for anaphylaxis management.

SCOPE

This policy applies to:

- all staff, including casual relief staff and volunteers
- all students who have been diagnosed with anaphylaxis, or who may require emergency treatment for an anaphylactic reaction, and their parents and carers.

POLICY

School Statement

Mornington Park Primary School will fully comply with Ministerial Order 706 and the associated guidelines published by the Department of Education and Training.

Anaphylaxis

Anaphylaxis is a severe allergic reaction that occurs after exposure to an allergen. The most common allergens for school-aged children are nuts, eggs, cow’s milk, fish, shellfish, wheat, soy, sesame, latex, certain insect stings and medication.

Symptoms

Signs and symptoms of a mild to moderate allergic reaction can include:

- swelling of the lips, face and eyes
- hives or welts
- tingling in the mouth.

Signs and symptoms of anaphylaxis, a severe allergic reaction, can include:

- difficult/noisy breathing
- swelling of tongue
- difficulty talking and/or hoarse voice
- wheeze or persistent cough
- persistent dizziness or collapse
- student appears pale or floppy
- abdominal pain and/or vomiting.

Symptoms usually develop within ten minutes and up to two hours after exposure to an allergen, but can appear within a few minutes.

Treatment

Adrenaline given as an injection into the muscle of the outer mid-thigh is the first aid treatment for anaphylaxis.

Individuals diagnosed as being at risk of anaphylaxis are prescribed an adrenaline autoinjector for use in an emergency. These adrenaline autoinjectors are designed so that anyone can use them in an emergency.

Individual Anaphylaxis Management Plans

All students at Mornington Park Primary School who are diagnosed by a medical practitioner as being at risk of suffering from an anaphylactic reaction must have an Individual Anaphylaxis Management Plan. When notified of an anaphylaxis diagnosis, the Principal of Mornington Park Primary School is responsible for developing a plan in consultation with the student's parents/carers.

Where necessary, an Individual Anaphylaxis Management Plan will be in place as soon as practicable after a student enrolls at Mornington Park Primary School and where possible, before the student's first day.

Parents and carers must:

- obtain an ASCIA Action Plan for Anaphylaxis from the student's medical practitioner and provide a copy to the school as soon as practicable
- immediately inform the school in writing if there is a relevant change in the student's medical condition and obtain an updated ASCIA Action Plan for Anaphylaxis
- provide an up-to-date photo of the student for the ASCIA Action Plan for Anaphylaxis when that Plan is provided to the school and each time it is reviewed
- provide the school with a current adrenaline autoinjector for the student that has not expired;
- participate in annual reviews of the student's Plan.

Each student's Individual Anaphylaxis Management Plan must include:

- information about the student's medical condition that relates to allergies and the potential for anaphylactic reaction, including the type of allergies the student has
- information about the signs or symptoms the student might exhibit in the event of an allergic reaction based on a written diagnosis from a medical practitioner
- strategies to minimise the risk of exposure to known allergens while the student is under the care or supervision of school staff, including in the school yard, at camps and excursions, or at special events conducted, organised or attended by the school
- the name of the person(s) responsible for implementing the risk minimisation strategies, which have been identified in the Plan
- information about where the student's medication will be stored
- the student's emergency contact details
- an up-to-date ASCIA Action Plan for Anaphylaxis completed by the student's medical practitioner.

Review and updates to Individual Anaphylaxis Management Plans

A student's Individual Anaphylaxis Management Plan will be reviewed and updated on an annual basis in consultation with the student's parents/carers. The plan will also be reviewed and, where necessary, updated in the following circumstances:

- as soon as practicable after the student has an anaphylactic reaction at school
- if the student's medical condition, insofar as it relates to allergy and the potential for anaphylactic reaction, changes
- when the student is participating in an off-site activity, including camps and excursions, or at special events including fetes and concerts.

Our school may also consider updating a student's Individual Anaphylaxis Management Plan if there is an identified and significant increase in the student's potential risk of exposure to allergens at school.

Location of plans and adrenaline autoinjectors

A copy of each student's Individual Anaphylaxis Management Plan will be stored with their ASCIA Action Plan for Anaphylaxis in all Staff Rooms, First Aid Room, Classrooms in folder in a prominent wall storage box. Individual student adrenaline autoinjectors will be kept in the student's classroom on a hook near the door with their ASCIA Action Plan. All staff and students know where these are.

Adrenaline autoinjectors for general use are available at First Aid Room, stored in a clearly labelled, bright yellow First Aid box.”.

Risk Minimisation Strategies

To reduce the risk of a student suffering from an anaphylactic reaction at Mornington Park Primary School, we have put in place the following strategies:

- *staff and students are regularly reminded to wash their hands after eating;*
- *students are discouraged from sharing food*
- *garbage bins at school are to remain covered with lids to reduce the risk of attracting insects*
- *gloves must be worn when picking up papers or rubbish in the playground;*
- *school canteen staff are trained in appropriate food handling to reduce the risk of cross-contamination*
- *year groups will be informed of allergens that must be avoided in advance of class parties, events or birthdays*
- *a general use EpiPen will be stored in the Sick Bay in the Main Building*
- *Planning for off-site activities will include risk minimisation strategies for students at risk of anaphylaxis including supervision requirements, appropriate number of trained staff, emergency response procedures and other risk controls appropriate to the activity and students attending.*
- *Ensure tables and bench tops are washed down after eating*
- *Ensure hand washing for all children upon arrival at the school, before and after eating*
- *Restrict use of food and food containers, boxes and packaging in crafts, cooking and science experiments, depending on the allergies of particular children*
- *Staff should discuss the use of foods in such activities with parents/carers of this child and these foods should be consistent with the risk minimisation plan.*
- *The risk minimisation plan will inform the school’s food purchases and menu planning.*

- *All children will be closely supervised at meal and snack times and consume food in specified areas. To minimise risk, children should not 'wander around' the school with food*
- *Staff should use non-food rewards, such as stickers, for all children*
- *Food preparation personnel (staff and volunteers) will be instructed about measures necessary to prevent cross contamination between foods during the handling, preparation and serving of food – such as careful cleaning of food preparation areas and utensils*
- *Where food is brought from home to the school, all parents/guardians/carers will be asked to avoid sending food containing specified allergens or ingredients as determined in the risk minimisation plan*

Adrenaline autoinjectors for general use

Mornington Park Primary School will maintain a supply of an adrenaline autoinjectors for general use, as a back-up to those provided by parents and carers for specific students, and also for students who may suffer from a first time reaction at school.

Adrenaline autoinjectors for general use will be stored at First Aid Room in a yellow labelled First Aid Box.

There are currently 4 adrenaline devices approved by the Therapeutic Goods Administration for use in Australia: the EpiPen[®], the Anapen[®], Jext[®] and Neffy[®]. All devices can be used when provided by families for students, however, the principal or allocated staff member can only use EpiPen[®], Anapen[®] or Jext[®] adrenaline autoinjector for general use. For more information about which autoinjector to purchase for general use, refer to Adrenaline autoinjectors for general use.

The Principal is responsible for arranging the purchase of adrenaline autoinjectors for general use, and will consider:

- the number of students enrolled at Mornington Park Primary School at risk of anaphylaxis
- the accessibility of adrenaline autoinjectors supplied by parents
- the availability of a sufficient supply of autoinjectors for general use in different locations at the school, as well as at camps, excursions and events
- the limited life span of adrenaline autoinjectors, and the need for general use adrenaline autoinjectors to be replaced when used or prior to expiry.
- the weight of the students at risk of anaphylaxis to determine the correct dosage of adrenaline autoinjector/s to purchase.

Emergency Response

In the event of an anaphylactic reaction, the emergency response procedures in this policy must be followed, together with the school's general first aid procedures, emergency response procedures and the student's Individual Anaphylaxis Management Plan.

A complete and up-to-date list of students identified as being at risk of anaphylaxis is maintained by the school's admin staff and stored at various locations including; Individual children's files, staff rooms, First Aid bags and alerts on Sentral. For camps, excursions and special events, a designated staff member will be responsible for maintaining a list of students at risk of anaphylaxis attending the special event, together with their Individual Anaphylaxis Management Plans and adrenaline autoinjectors, where appropriate.

If a student experiences an anaphylactic reaction at school or during a school activity, school staff must:

2.

Step	Action
1.	<ul style="list-style-type: none"> • Lay the person flat • Do not allow them to stand or walk • If breathing is difficult, allow them to sit with legs outstretched • Be calm and reassuring • Do not leave them alone • Seek assistance from another staff member or reliable student to locate the student's adrenaline device or the school's general use autoinjector, and the student's Individual Anaphylaxis Management Plan, stored in their classroom • If the student's plan is not immediately available, or they appear to be experiencing a first time reaction, follow steps 2 to 5
2.	<p data-bbox="331 1339 756 1370"><u>Administer an EpiPen or EpiPen Jr .</u></p> <ul style="list-style-type: none"> • Remove from plastic container • Form a fist around the EpiPen and pull off the blue safety release (cap) • Hold leg still and place orange end against the student's outer mid-thigh (with or without clothing) • Push down hard until a click is heard or felt and hold in place for 3 seconds · Remove EpiPen • Note the time the EpiPen is administered · Retain the used EpiPen to be handed to ambulance paramedics along with the time of administration. <p data-bbox="331 1729 743 1760"><u>OR Administer an Anapen® 500</u></p> <ul style="list-style-type: none"> • Pull off the black needle shield • Pull off grey safety cap (from the red button) • Place needle end firmly against the student's outer mid-thigh at 90 degrees (with or without clothing) • Press red button so it clicks and hold for 3 seconds • Remove Anapen® • Note the time the Anapen is administered

	<ul style="list-style-type: none"> Retain the used Anapen to be handed to ambulance paramedics along with the time of administration. <p><u>OR Administer Jext 150 or 300</u></p> <ul style="list-style-type: none"> Form fist around Jext and pull off yellow cap Place black injector tip against outer-mid thigh (with or without clothing) Push black tip firmly until a click is heard and hold in place for 3 seconds. Remove Jext Note the time the Jext device is administered. The used adrenaline device must be handed to the ambulance paramedics along with the time of administration <p><u>OR Administer Neffy® 1mg or 2mg</u></p> <ul style="list-style-type: none"> Hold the nasal spray with your thumb on the bottom of the plunger and a finger on either side of the nozzle. Do not pull or push on the plunger. Do not test or prime (pre-spray). Each Neffy nasal spray contains only one spray. Place the nozzle of the nasal spray into a nostril until fingers touch the nose. For smaller nostrils, aim for the fingers to touch the nose.
3.	Call an ambulance (000)
4.	If there is no improvement or severe symptoms progress (as described in the ASCIA Action Plan for Anaphylaxis), further adrenaline doses may be administered every five minutes, if other adrenaline autoinjectors are available.
5.	Contact the student's emergency contacts.

If a student appears to be having a severe allergic reaction, but has not been previously diagnosed with an allergy or being at risk of anaphylaxis, school staff should follow steps 2 – 5 as above.

Communication Plan

This policy will be publicly available on Mornington Park PS's website so that parents and other members of the school community can easily access information about Mornington Park PS's anaphylaxis management procedures.

The Principal is responsible for ensuring that all relevant staff, including casual relief staff, and volunteers are aware of this policy and procedures for anaphylaxis management. Casual relief staff and volunteers who are responsible for the care and/or supervision of students who are identified as being at risk of anaphylaxis will also receive a verbal briefing on this policy, their role in responding to an anaphylactic reaction and where required, the identity of students at risk.

The Principal will ensure that while students at risk of anaphylaxis are under the care or supervision of the school outside of normal class activities, including in the school yard, at camps and excursions, or at special event days, there is a sufficient number of school staff present who have been trained in anaphylaxis management.

The Principal is also responsible for ensuring relevant staff are trained and briefed in anaphylaxis management, consistent with the department's Anaphylaxis Guidelines.

Staff training

The Principal will ensure that the following school staff are appropriately trained in anaphylaxis management:

- School staff who conduct classes attended by students who are at risk of anaphylaxis
- School staff who conduct specialist classes, admin staff, first aiders and any other member of school staff as required by the Principal based on a risk assessment.

Staff who are required to undertake training must have completed:

- an approved face-to-face anaphylaxis management training course in the last three years, or
- an approved online anaphylaxis management training course in the last two years.

Example School uses the following training course [insert detail, for example ASCIA eTraining course (with 22579VIC), or 22578VIC.

Staff are also required to attend a briefing on anaphylaxis management and this policy at least twice per year (with the first briefing to be held at the beginning of the school year), facilitated by a staff member who has successfully completed an anaphylaxis management course within the last 2 years including School Anaphylaxis Supervisor.

Each briefing should address:

- the school's anaphylaxis management policy
- the causes, symptoms and treatment of anaphylaxis
- the identities of students diagnosed as being at risk of anaphylaxis, their allergens and the location of their Individual Anaphylaxis Management Plans and their medication/s
- discussion on staff anaphylaxis training and renewal
- how to use an adrenaline autoinjector, including hands-on practice with an adrenaline autoinjector trainer device (which does not contain adrenaline)
- the school's general first aid and emergency procedures
- the location of adrenaline autoinjector devices prescribed for individual students that have been purchased by their family
- the location of adrenaline autoinjector devices that the school has purchased for general use
- how to access on-going support and training.

When a new student enrolls at Mornington Park PS who is at risk of anaphylaxis, the Principal will develop an interim plan in consultation with the student's parents and ensure that appropriate staff are trained and briefed as soon as possible.

A record of staff training courses and briefings will be maintained and kept in school admin files and in the Emergency Management Plan.

FURTHER INFORMATION AND RESOURCES

- The Department's Policy and Advisory Library (PAL):
 - [Anaphylaxis](#)
 - [Allergy & Anaphylaxis Australia](#)
 - ASCIA Guidelines: [Schooling and childcare](#)
 - Royal Children's Hospital: [Allergy and immunology](#)

The Principal will complete the Department's Annual Risk Management Checklist for anaphylaxis management to assist with the evaluation and review of this policy and the support provided to students at risk of anaphylaxis.